

Shanghai Institute of Materia Medica¹, Chinese Academy of Sciences, Shanghai, and Guangzhou Institute of Chemistry², Chinese Academy of Sciences, Guangzhou, China

Synthesis and analgesic activity of stereoisomers of *cis*-fluoro-ohmefentanyl

ZOU YONG², WU HAO¹, YUAN WEIFANG¹, DAI QIYUAN¹, CHEN XINJIAN¹, JIN WENQIAO¹, ZHU YOUCHENG¹

Received August 22, 2002, accepted October 28, 2002

Dr. Zou Yong, Guangzhou Institute of Chemistry, Chinese Academy of Sciences, 1122 Mailbox, Guangzhou/Guangdong, P.R. China
zouyong@gic.ac.cn

Pharmazie 58: 300–302 (2003)

Four stereoisomers **1a–d** of *cis*-fluoro-ohmefentanyl have been synthesized. Their absolute configurations were determined by X-ray analysis of (3*S*,4*R*,2'*S*)-(–)-*cis*-**1d**. The analgesic activity (mice, sc, hot plate) revealed extreme stereodifferences. The ED₅₀ value of (3*R*,4*S*,2'*S*)-(+)-*cis*-**1a** was 0.000774 mg/kg (17958 times more potent than that of morphine), while the corresponding antipode **1c** was almost inactive.

1. Introduction


Ohmefentanyl (OMF) is a well-known potent μ -opioid receptor agonist which has firstly been synthesized in our laboratory in the early 1970s [1]. There are 3 chiral carbon atoms in OMF, so eight optically active isomers are possible. In our earlier reports [2, 3], we described the synthesis, stereochemistry, analgesic activity and opioid receptor binding characteristics of OMF. We have also found that the analgesic activity of the four stereoisomers of *cis*-OMF show great differences. For example, the ED₅₀ value (mice, ip, hot plate) of (+)-*cis*-(3*R*,4*S*,2'*S*)-OMF was 0.00106 mg/kg (13100 times more potent than that of morphine); while the enantiomer of which, (–)-*cis*-(3*S*,4*R*,2'*R*)-OMF was almost inactive under the same conditions (ED₅₀ > 10 mg/kg). At the same time, the analgesic activity of the four stereoisomers of *trans*-OMF

reveal little differences. Thus, it is believed that the *cis*-isomers of OMF can provide more information about the stereochemistry of the μ -opioid receptor, and be more useful for further research.

The above-mentioned results stimulated our interest in synthesizing and evaluating the four stereoisomers of *cis*-fluoro-ohmefentanyl (*cis*-FOMF). An F-atom was introduced in the aromatic ring mainly for two reasons, one is that the F-atom is an effective bioisoster known from many potent drugs; the other is that it is part of ¹⁸F substituted ohmefentanyl (¹⁸F-OMF), an extremely useful PET probe for molecular biology studies of the μ -opioid receptor.

Here we describe the synthesis of the four stereoisomers of *cis*-fluoro-ohmefentanyl, the determination of their absolute configurations, and the evaluation of their analgesic activities.

Scheme



Reagents: (a) NaBH₄/MeOH; (b) 10% Pd–C/H₂; (c) tartaric acid resolution; (d) 2-bromo-4'-fluoroacetophenone; (e) EtCOCl/PhMe; (f) NaBH₄/MeOH; (g) fractional crystallization.

choose ^{18}F -OMF as the PET probes in molecular pharmacology research.

(-)-*cis*-(3*R*,4*S*)-**6** (**6a**) was about 205 times more potent than morphine. We have also noticed that the duration of action maintaining above 50% of E_{max} was 300 min, much longer than that of other isomers which no more than 60 min, indicating the existence of a ketone may delay the metabolism of the compound. This result provided useful information for the research of long-acting analgesics.

4. Experimental

M.p.s. were determined with a Büchi 510 apparatus and are uncorrected. IR spectra were measured with a Perkin-Elmer 983G grating infrared spectrophotometer from KBr pellets. The ^1H NMR spectra were recorded on a Varian Genimi-2000 at 200 MHz using TMS as internal reference and CDCl_3 as solvent. MS spectra were recorded on a MAT-95 apparatus. Elemental analysis was performed on a Carlo-Erba 1106 apparatus. Optical rotation were measured on a Perkin-Elmer 241 polarimeter with a path length of 1 dm, concentrations are given in g/ml.

4.1. *cis*-1-Benzyl-3-methyl-*N*-phenyl-4-piperidinamine (**3**)

Compound **3** was prepared and separated according to the procedure described elsewhere [4].

4.2. (3*R*,4*S*)-3-Methyl-*N*-4-phenyl-4-piperidinamine (**4a**) and (3*S*,4*R*)-3-methyl-4-phenyl-4-piperidinamine (**4b**)

Preparation and optical resolution of **4a**, **4b** was performed according to the general procedure described elsewhere [3, 5].

4.3. (3*R*,4*S*)-*cis*-(+)-*N*-[1-[2-Hydroxy-2-(4'-fluorophenylethyl)]-4-piperidyl]-*N*-phenylpropanamide (**6a**) and (3*S*,4*R*)-*cis*-(-)-*N*-[3-methyl-1-[2-oxo-2-(4'-fluorophenylethyl)]-4-piperidyl]-*N*-phenylpropanamide (**6b**)

To a solution of **4a** (1.0 g) in dry toluene (25 ml) were added K_2CO_3 (8.0 g) and several crystals of KI. After stirring for 5 min, 2-bromo-4'-fluoroacetophenone (1.14 g) was added, the resultant mixture was stirred at room temperature for 1 h, filtered to remove solid, and obtaining the toluene solution of **5a**. As **5a** is rather unstable, the mixture was immediately acylated with propionyl chloride (2.0 ml) and refluxed for 5 h, allowed to cool, and then treated with aqueous K_2CO_3 and extracted with ether. The organic mixture was extracted with aqueous hydrochloride for several times, the combined acidic extracts were alkalized with K_2CO_3 extracted with ether, dried and concentrated. The residue was treated with EtOH saturated with HCl gas, recrystallization from EtOH/EtOAc/petroleum ether yielded white crystals, **6a**·HCl, m.p. = 236–238 °C, $[\alpha]_{\text{D}}^{25} - 6.96$ (c 0.39, MeOH). MS, m/z (%): 382 (M^+ 10), 335 (25) 259 (79), 160 (40), 122 (76), 105 (100), 77 (56).

6b·HCl was prepared from **4b** in the same procedure as **6a**·HCl, white crystals, m.p. 236–238 °C, $[\alpha]_{\text{D}}^{25} + 6.73$ (c 0.52, MeOH), MS, m/z (%): 382 (M^+ 10), 259 (100), 216 (20), 203 (38), 160 (36), 132 (9), 77 (5).

4.4. (3*R*,4*S*,2'*S*)-*cis*-(+)-*N*-[1-[2-Hydroxy-2-(4'-fluorophenylethyl)]-3-methyl-4-piperidyl]-*N*-phenyl-propanamide (**1a**) and (3*R*,4*S*,2'*R*)-*cis*-(-)-*N*-[1-[2-hydroxy-2-(4'-fluorophenylethyl)]-3-methyl-4-piperidyl]-*N*-phenylpropanamide (**1b**)

Solid NaBH_4 (0.5 g) was added in portions to a solution of **6a**·HCl (1.5 g) in MeOH (50 mL). The resultant mixture was refluxed for 2 h and then cooled, MeOH was evaporated with EtOAc several times, the EtOAc layer was dried and evaporated to obtain a mixture of **1a** and **1b**. Fractional crystallization from petroleum ether afforded the former crystallized **1a** (0.4 g) and the latter crystallized **1b** (0.35 g), respectively.

1a, fine white crystals, m.p. = 124–125 °C, $[\alpha]_{\text{D}}^{25} + 19.81$ (c 0.31, MeOH), ^1H NMR (CDCl_3) δ : 1.01 (3 H, t, $J = 7.4$ Hz, 10- CH_3), 1.17 (3 H, d, $J = 7.1$ Hz, 11- CH_3), 1.36 (1 H, br, 5e-H), 1.43 (1 H, br, 5a-H), 1.94 (2 H, q, $J = 7.6$ Hz, 9- CH_2), 2.11 (1 H, br, 6a-H), 2.38 (2 H, br, CH_2N), 2.66 (2 H, br, 2- CH_2), 2.76 (1 H, br, 3-CH), 3.04 (1 H, br, 6e-H), 3.99 (1 H, br, OH), 4.43 (1 H, dt, $J = 12.6$ Hz, 4a-H), 4.63 (1 H, br, 2'-CH), 6.97–7.9 (9 H, m, PhH). MS, m/z (%): 383 ($M-1$, 3), 366 (64), 323 (18), 259 (100), 216 (37), 203 (63), 160 (92), 149 (59), 132 (18), 77 (30).

1b, compact white crystals, m.p. = 133–135 °C, $[\alpha]_{\text{D}}^{25} - 29.78$ (c 0.45, MeOH), ^1H NMR (CDCl_3) δ : 1.01 (3 H, t, $J = 7.6$ Hz, 10- CH_3), 1.16 (3 H, d, $J = 7.1$ Hz, 11- CH_3), 1.35 (1 H, d, $J = 13.3$ Hz, 5e-H), 1.46 (1 H, qd, $J = 13.3$ Hz, 4.15 Hz, 5a-H), 1.93 (2 H, q, $J = 7.5$, 9- CH_2), 2.36 (4 H, m, 2a-H, 6a-H, 1'- CH_2), 2.69 (1 H, d, $J = 10.7$ Hz, 6e-H), 2.82 (1 H, br, 3e-H), 2.94 (1 H, d, $J = 11.5$ Hz, 2e-H), 4.44 (1 H, dt, $J = 12.9$ Hz, 4.34 Hz, 4a-H), 4.63 (1 H, dd, $J = 7.14$ Hz, 3.2 Hz, 2'-CH), 6.98–7.93 (9 H, m, PhH). MS, m/z (%): 385 ($M+1$, 4), 366 (8), 259 (100), 216 (30), 203 (58), 160 (52), 132 (10), 77 (10).

4.5. (3*S*,4*R*,2'*R*)-*cis*-(-)-*N*-[1-[2-Hydroxy-2-(4'-fluorophenylethyl)]-3-methyl-4-piperidyl]-*N*-phenyl-propanamide (**1c**) and (3*S*,4*R*,2'*S*)-*cis*-(+)-*N*-[1-[2-hydroxy-2-(4'-fluorophenylethyl)]-3-methyl-4-piperidyl]-*N*-phenylpropanamide (**1d**)

1c, **1d** were prepared and separated in the same procedure as **1a**, **1b**.

1c, fine white crystals, m.p. = 124–125 °C, $[\alpha]_{\text{D}}^{25} - 19.53$ (c 0.83, MeOH), ^1H NMR (CDCl_3) δ : 1.01 (3 H, t, $J = 7.4$ Hz, 10- CH_3), 1.16 (3 H, d, $J = 6.9$ Hz, 11- CH_3), 1.36 (1 H, br, 5e-H), 1.44 (1 H, d, $J = 9.1$ Hz, 5a-H), 1.94 (2 H, q, $J = 7.1$ Hz, 9- CH_2), 2.06 (1 H, mbr, 6a-H), 2.36 (2 H, m, 1'- CH_2N), 2.65 (2 H, br, 2- CH_2), 2.78 (1 H, br, 3-CH), 3.05 (1 H, d, $J = 11.0$ Hz, 6e-H), 4.05 (1 H, br, OH), 4.45 (1 H, dt, $J = 12.6$ Hz, 4.3 Hz, 4a-H), 4.61 (1 H, d, $J = 7.97$, 2'-CH), 6.97–7.39 (9 H, m, PhH). MS, m/z (%): 385 ($M+1$, 4), 366 (7), 259 (100), 216 (25), 203 (50), 160 (48), 132 (10), 77 (9).

1d, compact white crystals, m.p. = 133–135 °C, $[\alpha]_{\text{D}}^{25} + 27.77$ (c 0.66, MeOH), ^1H NMR (CDCl_3) δ : 1.01 (3 H, t, $J = 7.56$ Hz, 10- CH_3), 1.16 (3 H, d, $J = 6.87$ Hz, 11- CH_3), 1.34 (1 H, d, $J = 13.4$ Hz, 5e-H), 1.46 (1 H, qd, $J = 13.3$ Hz, 4.15 Hz, 5a-H), 1.92 (2 H, q, $J = 7.6$ Hz, 9- CH_2), 2.35 (4 H, m, 2a-H, 6a-H, 1'- CH_2), 2.68 (1 H, d, $J = 10.2$ Hz, 6e-H), 2.83 (1 H, br, 3e-H), 4.45 (1 H, dt, $J = 12.6$ Hz, 4.3 Hz, 4a-H), 4.61 (1 H, dd, $J = 10.2$ Hz, 2.7 Hz, 2'-CH), 6.98–7.40 (9 H, m, PhH). MS, m/z (%): 385 ($M+1$, 1), 366 (3), 259 (35), 238 (100), 223 (39), 203 (16), 160 (20), 146 (32), 77 (9).

Acknowledgements: This work was supported by The National Natural Science Foundation of China (Grant number: 29790123). The authors are very grateful to the staff of Department of Analysis, Shanghai Institute of Organic Chemistry, Chinese Academy of Science for the X-ray study.

References

- Jin, W. Q.; Xu, H.; Zhu, Y. C.; Fang, S. N.; Xia, X. L.; Huang, Z. M.; Ge, B. L.: Sci. Sin. (Engl. Ed.) **24**, 710 (1981)
- Wang, Z. X.; Zhu, Y. C.; Chen, X. J.; Ji, R. Y.: Chin. Sci. Bull. **15**, 1432 (1994)
- Wang, Z. X.; Zhu, Y. C.; Jin, W. Q.; Chen, X. J.; Chen, J.; Ji, R. Y.; Chi, Z. Q.: J. Med. Chem. **38**, 3652 (1995)
- Zou, Y.; Yuan, W. F.; Wu, H.; Dai, Q. Y.; Zhu, Y. C.: Chin. J. Pharm. **38**, 279 (1999)
- Van Bever, W. F. M.; Niemegeers, J. E.; Janssen, P. A.: J. Med. Chem. **17**, 1047 (1974)
- Brine, G. A.; Stark, P. A.; Liu, Y.; Carroll, F. I.; Singh, P.; Xu, H.; Rothman, R. B.: J. Med. Chem. **38**, 1547 (1995)
- Wang, Z. X.; Zhu, Y. C.; Chen, X. J.; Ji, R. Y.: Acta. Pharmaceutica Sinica. **28**, 950 (1993)